



Lecture 2.

Presheaf & Sheaf.



- Outline
- Presheaf, germ and stalk.
 - Sheafification.
 - Sheaf homomorphism $\left\{ \begin{array}{l} \text{injective} \\ \text{surjective} \\ \text{isomorphism} \\ \text{exact sequence} \end{array} \right.$
 - Examples.

- Throughout, X will be a topological space.
- Motivation: consider open sets $U \subseteq V \subseteq \mathbb{R}^n$.



f smooth on V , then certainly f is smooth on U .
 g is not smooth on V but it is smooth on U .

So the information of smooth functions on U is richer than on V .

• Def (Presheaf of group/ring/module)

A presheaf \mathcal{F} over X associates each open $U \subseteq X$ a group/ring/module $\mathcal{F}(U)$ s.t. the following holds:

① For $\forall V \subseteq U$, \exists a homomorphism

$$r_V^U : \mathcal{F}(U) \rightarrow \mathcal{F}(V)$$

s.t. $r_U^U = \text{id}$.

约定: $\mathcal{F}(\emptyset) = 0$.

② For $\forall W \subseteq V \subseteq U$ one has

$$r_W^V \circ r_V^U = r_W^U$$

• $\forall s \in \mathcal{F}(U)$ is called a section of \mathcal{F} on U .

• Example

put $r_V^U(s) := s|_V$.

① Constant sheaf. Let G be a group. Define a presheaf \mathcal{F} in the following way: for \forall connected open U ,


$$\mathcal{F}(U) := G.$$

$$\mathcal{F}_x = G \quad \forall x \in X$$

② Skyscraper: let $p \in X$. Let G be a group.

$$\mathcal{F}(U) := \begin{cases} G, & p \in U \\ 0, & p \notin U. \end{cases}$$


$$\mathcal{F}_x = \begin{cases} 0 & x \neq p \\ G & x = p \end{cases}$$



③ Two skyscrapers: Given $p \neq q \in X$.

Let G be a group.

$$F(U) := \begin{cases} G, & p \text{ or } q \in U \\ 0, & p \neq q \notin U. \end{cases}$$



But this is not sheaf.

$$F_x = \begin{cases} 0 & x \neq p \text{ or } q \\ G_1 & x = p \\ G_2 & x = q \end{cases}$$

④ presheaf of functions.

Let $X := \Omega \subseteq \mathbb{R}^n$ domain in \mathbb{R}^n .

For \forall open $U \subseteq \Omega$, let

$$F(U) := \left\{ \begin{array}{l} \text{continuous functions on } U \\ \uparrow \\ C^1, \text{ smooth, analytic, bounded.} \end{array} \right\}$$

⑤ presheaf of holomorphic/meromorphic functions.

Let $X := \Omega \subseteq \mathbb{C}^n$ be a domain in \mathbb{C}^n .

For \forall open $U \subseteq \Omega$, let

$$\mathcal{O}(U) := \{ \text{holo funct. on } U \}$$

$$\mathcal{O}_x = \left\{ \begin{array}{l} \text{convergent} \\ \text{power series} \\ \text{around } x \end{array} \right\}$$

$$\mathcal{M}(U) := \{ \text{meromorphic funct. on } U \}$$

$$\mathcal{O}^*(U) := \{ \text{holomorphic funct. w/ no zero on } U \}$$

$$\mathcal{B}(U) := \{ \text{bounded holomorphic function on } U \}$$

⑥ Ideal presheaf. Let $X := \mathbb{C}$.

$$I(U) := \begin{cases} \mathcal{O}(U), & 0 \notin U \\ \{ f \in \mathcal{O}(U) \mid f(0) = 0 \}, & 0 \in U \end{cases}$$

$$I_x = \mathcal{O}_x, \quad x \neq 0.$$

$I(U)$ is an ideal of $\mathcal{O}(U)$.

$$I_x = \{ f \in \mathcal{O}_x \mid f(0) = 0 \}$$

• Def A sheaf \mathcal{F} is a presheaf which satisfies the following two additional axioms

(A) If for $s, t \in \mathcal{F}(U) \exists$ open cover $U = \bigcup_{i \in I} U_i$ s.t. $s|_{U_i} = t|_{U_i}$ for $\forall i \in I$ then $s = t$.

(B) For open cover $U = \bigcup_{i \in I} U_i$ & $s_i \in \mathcal{F}(U_i)$ s.t. $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$ for $\forall i, j \in I$ $\exists s \in \mathcal{F}(U)$ s.t. $s|_{U_i} = s_i \forall i$.

▲ (A) means that \forall section is uniquely determined by its local information

(B) says that \forall compatible local sections glues together to a global section.

In the above examples, check that

③ two-scraper is not a sheaf.

⑤ \mathcal{B} is not a sheaf

• Germ & Stalk. Let \mathcal{F} be a presheaf.

Let $p \in X$ be a point. For \forall two open U & V both containing p


We say $s \in \mathcal{F}(U)$ & $t \in \mathcal{F}(V)$ have the same germ at p

if $\exists W \subseteq U \cap V$ containing p s.t. $s|_W = t|_W$.

In other words, a germ at p is an equivalence class

of sections: $\{s_i \in \mathcal{F}(U_i) \mid p \in U_i\} / \sim$ where

$$s_u \sim s_v \Leftrightarrow \exists_{p \in W} W \subseteq U \cap V \text{ s.t. } s_u|_W = s_v|_W.$$



e.g. $0 \in \mathbb{R} \subseteq \mathbb{C}^n$ connected.

For f & $g \in \mathcal{O}(\mathbb{R})$ w/ the same

germ at 0 , one must have $f = g$ on \mathbb{R}

In general, $f \in \mathcal{O}(U)$ & $g \in \mathcal{O}(V)$ have the same germ at $0 \in U \cap V$ iff f & g have the same power series expansion at 0 .

$\mathcal{F}_p := \{ \text{germ at } p \}$. This is called the stalk of \mathcal{F} at p .

▲ Note that for \forall open U containing p , there is a natural map $\mathcal{F}(U) \rightarrow \mathcal{F}_p$
 $s \mapsto s_p$.
 s_p is the germ of s at p .

• Ex. ① $\mathcal{O}_0 = \{ \text{convergent power series around } 0 \}$.
 ② Compute the stalks of all the previous examples.

Sheafification

Let \mathcal{F} be a presheaf. Then the sheafification of \mathcal{F} , denoted by \mathcal{F}^+ is given by

$$\mathcal{F}^+(U) := \left\{ \bigsqcup_{p \in U} \mathcal{F}_p \mid \text{for } \forall p \in U \exists p \in V \subseteq U \text{ \& } s \in \mathcal{F}(V) \text{ st. } s_p = s|_p, \forall p \in V \right\}$$


e.x. Show that \mathcal{F}^+ is a sheaf.

So there is a natural inclusion $\mathcal{F}(U) \hookrightarrow \mathcal{F}^+(U)$.

▲ e.x. When \mathcal{F} is a sheaf, then one actually has

$$\mathcal{F}(U) \xrightarrow{\cong} \mathcal{F}^+(U)$$

Example
 ▲ For two skyscrapers



$(U, 2)$ $(V, 3)$ cannot be glued in $\mathcal{F}(U \cup V)$

$(U, 2)$ & $(V, 3)$ give rise to a section in $\mathcal{F}^+(U \cup V)$

- In what follows, whenever we meet a presheaf, we will replace it by its sheafification, i.e., we will only deal w/ sheaves in the rest of this course.

- Sheaf morphisms. Let \mathcal{F} & \mathcal{G} be two sheaves

We say φ is a sheaf morphism from \mathcal{F} to \mathcal{G} if for \forall open U , \exists morphism

$\varphi_U: \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ s.t. the following diagram commutes for $\forall V \subseteq U$

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\varphi_U} & \mathcal{G}(U) \\ r_V^U \downarrow & & \downarrow r_V^U \\ \mathcal{F}(V) & \xrightarrow{\varphi_V} & \mathcal{G}(V) \end{array}$$

We call \mathcal{F} is a subsheaf of \mathcal{G} if φ_U is inclusion for all U .

- Given a sheaf morphism $\varphi: \mathcal{F} \rightarrow \mathcal{G}$.

① $\ker \varphi$ is given by $\ker \varphi(U) := \ker \varphi_U$.

e.x. $\ker \varphi$ is indeed a subsheaf of \mathcal{F} .

② $\text{Im} \varphi$ is the sheafification of the presheaf given by $\{\text{Im} \varphi_U\}_U$

- Given a subsheaf $\mathcal{F} \subseteq \mathcal{G}$,

\mathcal{G}/\mathcal{F} is the sheafification associated to $\left\{ \frac{\mathcal{G}(U)}{\mathcal{F}(U)} \right\}_U$

Def ① We say $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ injective if $\ker \varphi = 0$.

② We say $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ surjective if $\mathcal{G}/\text{Im} \varphi = 0$.

TFAE

- Prop: ① $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ is injective
- ② $\varphi_U: \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is injective for $\forall U$
- ③ $\varphi_x: \mathcal{F}_x \rightarrow \mathcal{G}_x$ is injective for $\forall x \in X$.

• Prop: TFAE

- ① $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ is surjective
- ② For $\forall \tau \in \mathcal{G}(U)$, \exists open cover $U = \cup U_i$
& $s_i \in \mathcal{F}(U_i)$ s.t. $\tau|_{U_i} = \varphi_{U_i}(s_i)$.
- ③ $\varphi_x: \mathcal{F}_x \rightarrow \mathcal{G}_x$ is surjective.

• We end this lecture by defining exactness.

let $\mathcal{F}, \mathcal{G}, \mathcal{Q}$ be sheaves.

Then $0 \rightarrow \mathcal{F} \xrightarrow{\alpha} \mathcal{G} \xrightarrow{\beta} \mathcal{Q} \rightarrow 0$ is called
an short exact sequence if

- ① α is injective
- ② β is surjective
- ③ $\ker \beta = \text{Im } \alpha$ (as sheaves)

• Prop. $0 \rightarrow \mathcal{F} \xrightarrow{\alpha} \mathcal{G} \xrightarrow{\beta} \mathcal{Q} \rightarrow 0$ is exact iff
 $0 \rightarrow \mathcal{F}_x \xrightarrow{\alpha_x} \mathcal{G}_x \xrightarrow{\beta_x} \mathcal{Q} \rightarrow 0$ is exact for $\forall x$.

* 讲义 : Coherent sheaves.